



LIVISTO
Along with you



GANADEXIL ENROFLOXACINA 5% 50 mg/ml

Solution for Injection
Enrofloxacin



- Effective at even low concentration
- $\geq 80\%$ Bioavailability
- Broad Spectrum
- Post - Antibiotic effect



Composition:

1 ml contains: Enrofloxacin 50mg

Description:

Ganadexil 5% Injection contains Enrofloxacin a broad spectrum antibiotic bactericidal in nature belongs to the group of Fluoroquinolones. Enrofloxacin inhibit the synthesis of bacterial DNA by interfering with bacterial enzyme action. This result in the death of bacteria. After injection blood concentration usually peaks in 1 hr. Bioavailability is more than 80%.

Antibacterial Spectrum:

Ganadexil 5% is active against a wide range of gram-negative organisms and several gram-positive bacteria. This includes E coli, Salmonella, Klebsiella, Enterobacter, Proteus, and generally Pseudomonas aeruginosa. It is also active against intracellular pathogens, including, eg, Brucella spp. also have significant activity against Mycoplasma and Chlamydia spp.

Indications:

Ganadexil 5% injection is indicated for the treatment of following bacterial infections in

Cattle, Calves, Buffalo, Sheep & Goat: for the treatment of Respiratory tract infections including Bronchopneumonia, Pasteurellosis and mycoplasmosis. Gastrointestinal tract infections including Diarrhea caused by colibacilli, colibacillosis and colisepticemia. also effective against mastitis and clostridial infections (enterotoxaemia).

Dogs & Cat: Treatment of: Respiratory, digestive, urinary and skin infections caused by microorganisms sensitive to enrofloxacin.

Dosage & Administration:

Cattle, Calves, Buffalo, Sheep & Goat: Subcutaneous use. 0.5-1 ml (equivalent to 25-50 mg of enrofloxacin)/10 kg of b.w.daily for 3 days. In the case of salmonellosis or severe infections, the length of treatment is 5 days. Do not apply more than 10 ml at the same injection site.

Dogs & Cats: subcutaneous use. 0.1 ml (equivalent to 5 mg of enrofloxacin)/kg of b.w daily for 5 days.

Withdrawal Period:

Meat: 14 days

Precautions:

Keep out of the sight and reach of children. Store below 25 °C. Protect from light.

Packing:

100ml Vial

علامات:

گینا ڈیکسل ایزوفلاکساسین گائے، بھینس، بھیڑ اور بکری میں سانس کے تمام جملہ امراض بشمول گل گھوٹو، نمونیہ، نظام انہضام کے امراض بشمول دست اور آنتوں کی سوزش اور انتڑیوں کے زہر اور ساڑو کا موثر علاج ہے۔

کتوں اور بلیوں میں سانس کے جراثیمی امراض، گردوں کے جراثیمی علاج میں موثر دوا ہے۔

خوراک و طریقہ استعمال:

گائے، بھینس، بھیڑ، بکری: 0.5-1 ملی لیٹر (بحساب 25-50 ملی گرام فی کلوگرام جسمانی وزن) انجکشن فی 10 کلوگرام جسمانی وزن تین دن تک کھال کے نیچے لگائیں۔
کتوں اور بلیوں میں 0.1 ملی لیٹر فی کلوگرام جسمانی وزن پانچ دن تک لگوائیں۔

بچوں کی پہنچ سے دور رکھیں، دھوپ سے بچائیں۔

احتیاط:

14 دن

وڈرل کا عرصہ: